



TAVI restricted to very old or very sick patients

Embargo Wednesday 29 August 08:30 CEST

Munich, Germany – August 29 2012: Cardiologists are restricting the use of transcatheter aortic valve implantation (TAVI) to very old or very sick patients at high surgical risk, according to research presented at the ESC Congress today. The first results of the Transcatheter Valve Treatment (TCVT) Sentinel Pilot Registry were presented by registry chairman Professor Carlo Di Mario.

The registry is part of the European Society of Cardiology (ESC) EURObservational Research Programme (EORP) of surveys and registries.

Today's presentation reveals current usage of the most modern TAVI valves and catheters in Europe, and compares indications, techniques and outcomes between different countries. "TAVI is a new technology which has been introduced in Europe but many question marks remain on which patients are most suitable," said Professor Di Mario. "We set up this registry because it was important to have a clear picture of clinical practice in Europe. Since our study was conducted during 2011-2012 we only included the very latest valves and delivery systems and this, together with the increased operator experience, probably explains the reduction in complications from previous studies and registries."

The registry included 4,571 patients who underwent the TAVI procedure using the Sapien XT or the CoreValve between January 2011 and May 2012. Patients were from 137 centres in Israel and 9 countries in Europe (Czech Republic, France, Spain, Switzerland, UK, Italy, Poland, Belgium, Germany).

The average age of patients was 81.4 ± 7.1 years, with equal numbers of men and women. There was a high prevalence of comorbidities in all patients, but patients who were 80 years old or younger had a greater incidence of diabetes, COPD, extracardiac arteriopathy (carotid, peripheral), permanent renal dialysis, previous myocardial infarction, previous cardiac surgery or percutaneous coronary intervention (PCI), previous aortic valve replacement (valve-in-valve procedure). Professor Di Mario said: "This shows that the use of TAVI in younger patients has been restricted to those with more comorbidities, who therefore have high surgical risks."

Overall in-hospital mortality was 7.4%. There were no significant differences in in-hospital mortality based on valve type (6.7% CoreValve, 7.9% Sapien XT, $p=0.15$) but there were significant differences based on the approach site (transfemoral 5.9%, transapical 12.8%, trans-subclavian and other approaches 9.7%, $p<0.01$).

The most frequent complication was bleeding requiring blood transfusion(s). The incidence was greater than 20% for the transapical and trans-subclavian approaches and 15.0% for the transfemoral approach ($p<0.01$).

The need to implant a permanent pacemaker was another frequent complication, with an overall incidence of 13.2%. The incidence of this complication was significantly greater in patients who received the CoreValve (23.4%) than in patients who received the Sapien XT valve (6.0%) ($p<0.01$).

There were no other differences in the incidence of in-hospital complications between the two valve types. There was a low and similar incidence of major vascular complications (3.1%), stroke (1.8%), myocardial infarction (0.9%), and other complications.

The length of hospital stay was 9.3 ± 8.1 days. There was wide variation between countries, from 7 days in Switzerland to more than 12 days in Poland. Patients who received general anaesthesia had longer hospital stays than patients who received local anaesthesia (10.2 ± 8.7 days vs 7.9 ± 6.1 days, $p < 0.01$). Hospital stays of more than 10 days occurred in 43.8% of patients treated with a transapical approach, 39.5% treated with other surgical approaches and 22.0% of patients treated transfemorally.

“Anaesthesia has a big influence on cost, duration of the procedure and length of hospital stay,” said Professor Di Mario. “There are big differences between countries in the type of anaesthesia used. Poland and the UK conduct most transfemoral procedures under general anaesthesia while Switzerland and Italy use local anaesthesia for more than 60% of these procedures.”

Pharmacological treatment varied greatly between countries. There was relatively low use of aspirin (64.3%) and clopidogrel or other thienopyridines (32.6%) at discharge. Professor Di Mario said: “This finding suggests poor compliance with the recommended strategy of using a combination of the two drugs at discharge after TAVI.”

He added: “With an in-hospital mortality of 7.3%, this contemporary registry of the most up-to-date valves and delivery catheters shows that TAVI remains a high risk procedure when applied to very old patients or patients with significant comorbidities who are poor surgical candidates.”

Professor Di Mario continued: “We now have a clear understanding of how TAVI is being used in these 10 countries. In the majority of cases, patients are very old and very sick, with lots of comorbidities which would have made their journey to surgery a nightmare.”

He concluded: “The next step we are considering will be to launch a permanent registry that involves all countries in Europe. Collecting information on how devices are used, whether they are used appropriately, and how practice varies between countries offers enormous opportunities to provide data to regulatory authorities, monitor adherence to best practice guidelines and improve care for patients.”

ENDS

Notes to Editor

About the [European Society of Cardiology](#)

The European Society of Cardiology (ESC) represents more than 75,000 cardiology professionals across Europe and the Mediterranean. Its mission is to reduce the burden of cardiovascular disease in Europe.

About ESC Congress 2012

The ESC Congress is currently the world's premier conference on the science, management and prevention of cardiovascular disease. ESC Congress 2012 takes place 25-29 August at the Messe München in Munich. The scientific programme is available [here](#).

More information is available from the ESC Press Office at press@escardio.org

REFERS TO SESSION: Clinical Trial & Registry Update III: Updates on Atrial Fibrillation and Valves, Webcast – Munich – Central Village – Wednesday 29 Aug 08:30-10:00
Session details available [here](#)